# Understanding Spatial and Temporal Variations of Arctic Circulation Using Oxygen Isotopes of Seawater


Undergraduate: Leon Yin
High School Student: Cassandra Kopans-Johnson
High School Teacher: Susan Meabh Kelly
Researcher: Allegra LeGrande







# **Purpose of Climate**

- Earth has a heat budget.
- As solar radiation bombards it, equatorial regions absorb more heat while polar regions absorb less heat.
- Earth redistributes heat through atmospheric and oceanic circulation.



https://www.e-education.psu.edu/files/earth540/image/Lesson4/HeatRedist.png






#### **Water Masses**

Regional pockets are surface sources for water mass formation.


They trigger thermohaline circulation, density driven oceanic circulation.



# Defining δ<sup>18</sup>O

- <sup>18</sup>O =
  - Heavy isotope of oxygen
  - A semi-conserved tracer of the hydrologic cycle due to its regional relationship with salinity
- In standard ocean water (R<sub>standard</sub>),
   For each 1,000,000 H<sub>2</sub><sup>16</sup>O molecules
   ~2005 H<sub>2</sub><sup>18</sup>O molecules
- 'delta' notation:  $\delta^{18}$ O in permil (%)  $\equiv (R_{\text{sample}}/R_{\text{standard}} - 1) X1000$

# Rayleigh Distillation



<sup>18</sup>O (heavy)

hard to evaporate, easy to precipitate

→ salt water

<sup>16</sup>O (light)

easy to evaporate, hard to precipitate

→ fresh water

## Fairbanks and Charles 1992




Fig 30.3B. Plots of  $\delta^{18}$ O and salinity measurements across all major salinity gradients in the Atlantic and eastern tropical Pacific (Fairbanks, ms in preparation)

#### **Research Context**

Scientists have been using <sup>18</sup>O water isotopologues as tracers of the hydrologic cycle to better understand ocean circulation.

#### **Previous Work**


- Database: (1999) G.
   Schmidt et al
- Gridded dataset: (2006)
   A. LeGrande and G.
   Schmidt

#### **Our Work**


- Update the database.
- Examine spatial and temporal impact on <sup>18</sup>O in the Arctic.
- Improve regional <sup>18</sup>O-S relationships to redefine Arctic water masses


Calculated Surface δ<sup>18</sup> O Seawater

Estimated Surface  $\delta^{18}$  O Seawater



 $\delta^{18}$  O Est-Calc Discrepencies







A. LeGrande and G. Schmidt and this project's updates


Calculated Surface δ<sup>18</sup> O Seawater

Estimated Surface  $\delta^{18}$  O Seawater





 $\delta^{18}$ O Est-Calc Discrepencies





A. LeGrande and G. Schmidt and this project's updates





#### **Spatial Distribution**


Coastal water - Open water




River runoff

=


Rayleigh Distillation and resulting inland freshwater discharge with lower  $\delta^{18}$ O.





#### **Temporal Distribution**



<sup>18</sup>O Levels







## **Define Arctic Water Masses**


#### Nature -


where does water originate from?


#### **Nurture** -


where has it been?


Unique qualities.














## Statistics for outliers



## **Redefined Water Masses**





# **Summary**

#### Goal:

To better understand the Arctic Ocean's complex circulation.

#### **Accomplishments:**

- Identified estimated-calculated discrepancies.
- Examined regional temporal and spatial distribution to define water masses.
- Used Jackknife to statistically analyze the water masses <sup>18</sup>O-S relationships.
- Consolidated ten water masses into four water masses.

#### **Future Work**

- Application of Jackknife slope analysis to improve regional  $\delta^{18}$ O-S relationships that better define water masses and track their sources, pathways, and interactions in global ocean circulation.
- Account for temporal variation in the Global Gridded Data Set.
- Representation of ocean components along isopycnals.
- Collect additional oxygen-18 data by reviewing non-English language scientific publications

#### References

Bauch, D., et al. (2010), Changes in distribution of brine waters on the Laptev Sea shelf in 2007. Journal of Geophysical Research 115, C11.

Bauch, D., P. Schlosser, and R. Fairbanks (1995), Freshwater balance and the source of deep and bottom waters in the Arctic Ocean inferred from the distribution of H218O. *Progress in Oceanography 35*: 53-80.

Craig, H., and L. I. Gordon (1965), Deuterium and oxygen 18 variations in the ocean and the marine atmosphere. In E. Tongiori (Ed.) *Proceedings of a Conference on Stable Isotopes in Oceanographic Studies and Paleotemperatures*, Spoleto, Italy.

Dansgaard, W. (1964), Stable isotopes in precipitation. Tellus 16, 436-468.

Grabs, W. E., F. Portmann, and T. De Couet (2000), Discharge Observation Networks in Arctic Regions: Computation of the River Runoff into the Arctic Ocean, Its Seasonality and Variability. In Lewis, E.L., Jones, E.P., Lemke, P., Prowse, T.D., and P. Wadhams (Eds.), *Freshwater Budget of the Arctic Ocean*. Boston: Kluwer Academic Publishers, 249-65.

Létolle, R., et al. (1993), <sup>18</sup>O abundance and dissolved silicate in the Lena delta and Laptev Sea (Russia). *Marine Chemistry 43*, 47-64.

LeGrande, A.N., and G. A. Schmidt (2006), Global gridded data set of the oxygen isotopic composition in seawater. Geophysical Research Letters 33.

Schmidt, G. A., G. R. Bigg, and E. J. Rohling (1999), Global seawater oxygen-18 database - v1.21. http://data.giss.nasa.gov/o18data/

# Acknowledgements

Allegra LeGrande Susan Meabh Kelly **Gavin Schmidt** Matthew Pearce Colleagues at GISS **NYCRI NASA GISS** NASA GSFC